red nose saving little lives

fast facts 2017



births in australia 2015¹ 305,377

6

money invested in research to date by red nose \$16.7 million



research currently being funded by red nose

- An individual participant data meta-analysis of going-tosleep position and risk of late pregnancy stillbirth
- The effect of migration and acculturation on risk of stillbirth in Western Australia
- Preventing Term Stillbirth in South Asian Born Mothers: A stepped wedge cluster randomised controlled trial



- definitions and explanations
- SUDI Sudden Unexpected Deaths in Infancy including SIDS.
- Perinatal All fetal and neonatal deaths of at least 400 grams birthweight or at least 20 weeks gestation. Deaths within 28 days of birth. Neonatal

Stillbirth Fetal deaths (if the birthweight is at least 400 grams or the gestational age is 20 weeks or more). Injury and poisoning deaths include persons aged 0-4 years in Australia per annum on average calculated using 2011-2014 data. See note 5 for source.

- 2 Australian Institute of Health and Welfare (2016). Australia's mothers and babies 2014-in brief. Perinatal statistics series no. 31. Cat no. PER 72. Canberra: AIHW.
- 3 Australian Bureau of Statistics (2016). 3303.0 Causes of Death, Australia, 2015. ABS, Canberra
- 4 Explanatory note from the ABS: "Data with small values have been randomly assigned to protect the confidentiality of individuals. As a result, some totals will not equal the sum of their components"
- 5 Australian Institute of Health and Welfare (2017). GRIM (General Record of Incidence of Mortality) Books 2014: All external causes of morbidity and mortality. Canberra: AIHW. Note: Data for all external causes of mortality are from the ICD-10 (V01-Y98).
- 6 Australian Bureau of Statistics (2003). SIDS in Australia 1981-2000: A statistical overview. ABS, Canberra & Australian Bureau of Statistics (2001-).
- 3303.0 Causes of Death, Australia, 2001-, ABS, Canberra 7 Includes calculations prepared by Red Nose and confirmed by the ABS